arXiv:1302.2133v1 [math.GT] 8 Feb 2013

Cubulated moves and discrete knots[*]

Gabriela Hinojosa[†] Alberto Verjosvky, [‡]
Cynthia Verjovsky Marcotte

January 21, 2013

Abstract

In this paper, we prove than given two cubic knots K_1, K_2 in $\mathbb{R}^3$, they are isotopic if and only if one can pass from one to the other by a finite sequence of cubulated moves. These moves are analogous to the Reidemeister moves for classical tame knots. We use the fact that a cubic knot is determined by a cyclic permutation of contiguous vertices of the $\mathbb{Z}^3$-lattice (with some restrictions), to describe some of the classic invariants and properties of the knots in terms of such cyclic permutations, by projecting onto a plane such that it is injective when restricted to the $\mathbb{Z}^3$-lattice and the image of the $\mathbb{Z}^3$-lattice is dense.

1 Introduction

In [3] it was shown that any smooth knot $\mathbb{S}^n \sim K^n \subset \mathbb{R}^{n+2}$ can be deformed isotopically into the n-skeleton of the canonical cubulation of $\mathbb{R}^3$. In particular, every classical smooth knot $\mathbb{S}^1 \subset \mathbb{R}^3$ is isotopic to a cubic knot. There are two elementary "cubulated moves". The first one (M1) is obtained by dividing each cube of the original cubulation into m^3 cubes, which means that each edge of the knot is subdivided into m equal segments. The second one (M2) consists in exchanging a connected set edges in a face of the cubulation (or a subdivision of the cubulation) with the complementary edges in that face. If two cubic knots K_1 and K_2 are such that we can convert K_1 to K_2 using a finite sequence of cubulated moves

---

[*] *2010 Mathematics Subject Classification.* Primary: 57M25. Secondary: 57M27, 57Q45. *Key Words.* Cubic knots, discrete knots.

[†] This work was partially supported by CONACyT (Mexico), CB-2009-129939

[‡] This work was partially supported by CONACyT (Mexico), CB-2009-129280 and and PAPIIT (Universidad Nacional Autónoma de México) #IN100811.

then we say that they are equivalent via cubulated moves, notated $K_1 \overset{c}{\sim} K_2$.

This allows us to prove the following:

Theorem 1. *Given two cubic knots K_1 and K_2 in $\mathbb{R}^3$, then they are isotopic if and only if K_1 is equivalent to K_2 by a finite sequence of cubulated moves;* i.e.*, $K_1 \overset{c}{\sim} K_2$.*

Theorem 1 is analogous to the Reidemeister moves of classical tame knots for cubic knots.

The fact that every knot is isotopic to a cubic knot allows us to develop a discrete version of a knot. Since a cubic knot is given by a sequence of edges whose boundaries are in the canonical lattice of points with integer coefficients in $\mathbb{R}^3$, *i*.e., the abelian group $\mathbb{Z}^3$, each knot is determined by a cyclic permutation $(a_1, \ldots, a_n)$ (with some restrictions), $a_i \in \mathbb{Z}^3$. We can describe some of the invariants and properties of the knots in terms of such cyclic permutations by projecting onto a plane P, such that it is injective when restricted to the $\mathbb{Z}^3$-lattice and the image of the $\mathbb{Z}^3$-lattice, Λ_P, is dense. More precisely, the projection of each knot is determined by a cyclic permutation $(w_1, \ldots, w_n)$ (with some restrictions), $w_i \in \Lambda_P$. In Section 5, we give a discrete version of Seifert surfaces and the Alexander polynomial. We also describe the fundamental group using the Wirtinger presentation.

2 Cubulations for $\mathbb{R}^{n+2}$

A *cubulation* of $\mathbb{R}^{n+2}$ is a decomposition of $\mathbb{R}^{n+2}$ into a collection $\mathcal{C}$ of $(n+2)$-dimensional hypercubes such that any two of its hypercubes are either disjoint or meet in one common face of dimension $k \leq n+1$. This provides $\mathbb{R}^{n+2}$ with the structure of a cubic complex.

In general, the category of cubic complexes and cubic maps is similar to the simplicial category. The main difference consists in using cubes instead of simplexes. In this context, a cubulation of a manifold is specified by a cubical complex PL-homeomorphic to the manifold (see [5], [8], [10]).

The *canonical cubulation* $\mathcal{C}$ of $\mathbb{R}^{n+2}$ is the decomposition of $\mathbb{R}^{n+2}$ into hypercubes which are the images of the unit cube $I^{n+2} = \{(x_1, \ldots, x_{n+2}) \,|\, 0 \leq x_i \leq 1\}$ by translations by vectors with integer coefficients.

Definition 2.1. The n-skeleton of $\mathcal{C}$, denoted by $\mathcal{S}$, consists of the union of the n-skeletons of the cubes in $\mathcal{C}$, *i.e.*, the union of all cubes of dimension n contained in the faces of the $(n+2)$-cubes in $\mathcal{C}$. We will call $\mathcal{S}$ the *canonical scaffolding* of $\mathbb{R}^{n+2}$.

Any cubulation of $\mathbb{R}^{n+2}$ is obtained from the canonical cubulation by applying to the canonical cubulation a conformal transformation $x \mapsto \lambda A(x) + a$ where $\lambda \neq 0$ is a real number, $a \in \mathbb{R}^{n+2}$ and $A \in SO(n+2)$.

Consider the homothetic transformation $\mathfrak{h}_m : \mathbb{R}^{n+2} \to \mathbb{R}^{n+2}$ given by $\mathfrak{h}_m(x) = \frac{1}{m}x$, where $m > 1$ is an integer. The set $\mathcal{C}_{\frac{1}{m}} := \mathfrak{h}_m(\mathcal{C})$ is called a *subcubulation* or *cubical subdivision* of $\mathcal{C}$.

In [3], we proved the following theorem which is central to this paper.

Theorem 2.2. *Let $\mathcal{C}$ be the standard cubulation of $\mathbb{R}^{n+2}$. Let $K \subset \mathbb{R}^{n+2}$ be a smooth knot of dimension n. There exists a knot $\hat{K}$ continuously isotopic to K, which is contained in the scaffolding $\mathcal{S}$ (n-skeleton) of the standard cubulation $\mathcal{C}$ of $\mathbb{R}^{n+2}$. The cubulation of the knot $\hat{K}$ admits a subdivision by simplexes and with this structure the knot is PL-equivalent to the n-sphere with its canonical PL-structure.*

3 Cubulated isotopy

Recall that a smooth parametrized knot is a smooth embedding $k : \mathbb{S}^n \to \mathbb{R}^{n+2}$. As is usual, we will at times identify the embedding with its image, which we call a geometrical knot.

Given two smooth parametrized n-dimensional knots K_1, $K_2 : \mathbb{S}^n \hookrightarrow \mathbb{R}^{n+2}$ we say they are smoothly isotopic if there exists a smooth isotopy $H : \mathbb{S}^n \times \mathbb{R} \to \mathbb{R}^{n+2}$ such that

$$H(x, \{t\}) = \begin{cases} K_1(x) & \text{if } t \leq -1, \\ K_2(x) & \text{if } t \geq 1, \end{cases}$$

and $H(\cdot, \{t\})$ is an embedding of $\mathbb{S}^n$ for $t \in \mathbb{R}$.

Definition 3.1. We will say $J = \{(H(x,t), t) \in \mathbb{R}^{n+3} \,|\, x \in \mathbb{S}^n \ t \in \mathbb{R}\}$ is the *isotopy cylinder* of K_1 and K_2. Note that J is a smooth submanifold of codimension two in $\mathbb{R}^{n+3}$.

Let $\mathcal{C}$ be the canonical cubulation of $\mathbb{R}^{n+3}$. Let $p : \mathbb{R}^{n+3} \hookrightarrow \mathbb{R}$ be the projection onto the last coordinate.

Definition 3.2. Let M be a connected subset of $\mathbb{R}^{n+3}$ such that $p^{-1}(c) \cap M$ or $p|_M^{-1}(c)$ is connected for all $c \in \mathbb{R}$, we say that M is *sliced by connected level sets of* p.

Observe that there is no restriction on the dimension of M.

Note that $p|_J$ is sliced by connected sets.

In this section we will prove:

Theorem 3.3. *The isotopy cylinder J can be cubulated. In other words, there exists an isotopic copy J' of J contained in the $(n+1)$-skeleton of the canonical cubulation of $\mathbb{R}^{n+3}$. Moreover J' can be chosen to be sliced by connected level sets of p.*

To prove Theorem 3.3 we need some preliminary results:

Proposition 3.4. *Let M be a closed subset of $\mathbb{R}^{n+3}$, such that $p|_M$ is a proper function and $p|_M^{-1}(c)$ is non-empty for all $c \in \mathbb{R}$. Assume M is sliced by connected level sets of p. Let $\mathcal{Q} \subset \mathcal{C}$ be the union of all cubes of the canonical cubulation intersecting M. Then $\mathcal{Q}$ is also sliced by connected level sets of p.*

Proof. Let c be any number in $\mathbb{R}$. Then c belongs to the closed interval $[n, n+1]$ for some number $n \in \mathbb{Z}$.

Let A be the intersection $p^{-1}[n, n+1] \cap M$, so A is compact in $\mathbb{R}^3$ and is sliced by connected level sets. Assume A is not connected, so $A = A_1 \cup A_2$ where A_1, A_2 are non-empty, disjoint compact subsets and both A_1 and A_2 are also sliced by connected level sets of p. Then $p(A_1) \cup p(A_2) = [n, n+1]$, $p(A_1) \cap p(A_2) = \emptyset$ and $p(A_1)$ and $p(A_2)$ are compact sets, hence $[n, n+1]$ is not connected, which is a contradiction. Therefore A is connected.

Claim 1: $p^{-1}[n, n+1] \cap \mathcal{Q}$ is connected.
Suppose that $p^{-1}[n, n+1] \cap \mathcal{Q} = C_1 \cup C_2$, where C_1, C_2 are non-empty disjoint closed sets. As A is connected, we have that either $A \subset C_1$ or $A \subset C_2$. We assume that $A \subset C_1$. But A intersects each of the hypercubes belonging to $p^{-1}[n, n+1] \cap \mathcal{Q}$, since each hypercube is connected, it follows that $p^{-1}[n, n+1] \cap \mathcal{Q} \subset C_1$, which is a contradiction. Therefore $p^{-1}[n, n+1] \cap \mathcal{Q}$

is connected.

Claim 2: $p|_{\mathcal{Q}}$ is sliced by connected level sets of p.
This is a consequence of the above claim and the fact that the cubes intersecting $p^{-1}(c)$ and $p^{-1}[n, n+1]$ are the same cubes, since $\mathcal{C}$ is the canonical cubulation. □

Recall that $K_1,\ K_2 \subset \mathbb{R}^{n+2}$ are two smoothly isotopic knots in $\mathbb{R}^{n+2}$ with $H : \mathbb{S}^n \times \mathbb{R} \to \mathbb{R}^{n+2}$ the smooth isotropy between them, which gives us the isotopy cylinder $J = \{(H(x,t), t) \in \mathbb{R}^{n+3} \,|\, x \in K_1\}$.

Lemma 3.5. *The isotopy cylinder J is sliced by connected level sets of p.*

Proof. Since $p^{-1}(t) = H(\cdot, t)$ is the image of an embedding of $\mathbb{S}^n$, and hence connected, the result follows. □

Lemma 3.6. *Let $\mathcal{Q}_{\mathcal{J}}$ be the union of all cubes which intersect J. Then $\mathcal{Q}_{\mathcal{J}}$ is sliced by connected level sets of p.*

Proof. Note that $p|_J$ is clearly proper, therefore this is a consequence of Proposition 3.4. □

Proof of Theorem 3.3. To finish the proof we only need to show that J can be cubulated by a small isotopy, in other words: there exists a small isotopy which moves J onto an isotopic copy of J embedded in the $n+1$-skeleton of a sufficiently small subdivision of the canonical cubulation of $\mathbb{R}^{n+3}$. But this is proven in [3]. □

4 Cubulated moves

We say that a knot $K \subset \mathbb{R}^3$ is a *cubic knot*, if K is contained in the scaffolding $\mathcal{S}$ of the canonical cubulation $\mathcal{C}$ of $\mathbb{R}^3$.

Definition 4.1. The following are the allowed *cubulated moves*:

M1 *Subdivision:* Given an integer $m > 1$, consider the subcubulation $\mathcal{C}_m$ of $\mathcal{C}$. Since $\mathcal{C} \subset \mathcal{C}_m$, then K is contained in the scaffolding $\mathcal{S}_m$ (the 1-skeleton) of $\mathcal{C}_m$ and as a cubic complex, each edge of K is subdivided into m equal edges.

M2 *Face Boundary Move:* Suppose that K is contained in some subcubulation $\mathcal{C}_m$ of the canonical cubulation $\mathcal{C}$ of $\mathbb{R}^3$. Let $Q \in \mathcal{C}_m$ be a cube such that $A = K \cap Q$ contains an edge. We can assume, up to applying the elementary move $M1$, that A consists of either one, two or three edges that are connected and are part of the boundary of a face $F \subset Q$. Thus A is an arc contained in the boundary of F and ∂F is divided by ∂A into two cubulated arcs. One of them is A and we denote the other by A'. Observe that both arcs share a common boundary. The move consists in replacing A by A'.

Remark 4.2. Notice that the face boundary move can be extended to an ambient isotopy of $\mathbb{R}^3$.

Definition 4.3. Given two cubic knots K_1 and K_2 in $\mathbb{R}^3$. We say that K_1 is *equivalent* to K_2 *by cubulated moves*, denoted by $K_1 \overset{c}{\sim} K_2$, if we can transform K_1 to K_2 by a finite number of cubulated moves.

4.1 Main theorem

We are now ready to prove Theorem 1.

Theorem 1. *Given two cubic knots K_1 and K_2 in $\mathbb{R}^3$, then they are isotopic if and only if K_1 is equivalent to K_2 by cubulated moves;* i.e., *$K_1 \overset{c}{\sim} K_2$.*

Proof. First, note that if K_1 and K_2 are equivalent by cubulated moves, then these knots must clearly be isotopic. Hence, what remains to be proved is that two cubic knots that are isotopic must also be equivalent by cubulated moves.

Our strategy is as follows. First, for $i \in \{1, 2\}$, we will smooth each K_i to obtain $\widetilde{K}_i$, and then cubulate these two knots to obtain K'_i in such a way that a) $K_i \overset{c}{\sim} K'_i$ and b) we can show $K'_1 \overset{c}{\sim} K'_2$ are equivalent by cubulated moves.

Given a cubic knot K, there exists a smooth knot $\widetilde{K}$ isotopic to K such that $\widetilde{K}$ is $\mathcal{C}^0$-arbitrarily close to K. This is because we can round the corners at the vertices of K in an arbitrarily small neighborhoods of them (see [6]).

Let J be the isotopy cylinder (defined above) of $\widetilde{K}_1$ and $\widetilde{K}_2$. Then J is a smooth submanifold of codimension two in $\mathbb{R}^4$. By Theorem 3.3, there exists an isotopic copy of J, say J', contained in the 2-skeleton of the canonical cubulation $\mathcal{C}$ of $\mathbb{R}^4$. Recall that J' is sliced by connected level sets of p. Furthermore, note that there exist integer numbers m_1 and m_2 such that

$p^{-1}(t) \cap J' = K_1'$ for all $t \leq m_1$ and $p^{-1}(t) \cap J' = K_2'$ for all $t \geq m_2$, where K_1' and K_2' are cubic knots which are isotopic to $\widetilde{K}_1$ and $\widetilde{K}_2$, respectively.

Now, we will use the following results which will be proved in Sections 4.1.1 and 4.1.2, respectively.

Lemma 4.4. *Given a cubic knot K we can choose a cubulation $\mathcal{C}_{\frac{1}{m}}$ fine enough that $N(K) = \cup\{Q \in \mathcal{C}_{\frac{1}{m}} \mid Q \cap K \neq \emptyset\}$ is a closed tubular neighborhood of K and $Q \cap K$ is equal to either vertex, one edge, two edges sharing a vertex. We can also choose $\widetilde{K}$ isotopic to K such that $\widetilde{K}$ is $\mathcal{C}^0$-arbitrarily close to K and $\widetilde{K} \subset Int(N(K))$. Then using Theorem 3.3 we obtain K' in $\partial N(K)$ cubic in $\mathcal{C}_{\frac{1}{m}}$, and $K \overset{c}{\sim} K'$;* i.e., *we can go from K to K' by cubulated moves.*

Theorem 4.5. *Given two cubic knots K_1 and K_2, we obtain K_1' and K_2' as in Lemma 4.4. Then there exists a finite sequence of cubulated moves that carries K_1' into K_2'. In other words, K_1' is equivalent to K_2' by cubulated moves.*

Thus by Lemma 4.4, there exists a finite sequence of cubulated moves that carries K_1 into K_1' and also a finite sequence of cubulated moves that carries K_2 into K_2'. By Theorem 4.5, there exists a finite sequence of cubulated moves that carries K_1' into K_2'. As a consequence there exists a finite sequence of cubulated moves that carries K_1 into K_2. $\square$

4.1.1 Cubic case

Let K be a cubic knot. We can choose a cubulation $\mathcal{C}_{\frac{1}{m}}$ fine enough such that the subcollection $\mathcal{N}_K = \{Q \in \mathcal{C}_{\frac{1}{m}} \mid Q \cap K \neq \emptyset\}$ satisfies that $N(K) = \cup_{Q \in \mathcal{N}_K} Q$ is a closed tubular neighborhood of K and $Q \cap K$ is equal to either vertex, one edge, two edges sharing a vertex. We can also choose $\widetilde{K}$ isotopic to K such that $\widetilde{K}$ is $\mathcal{C}^0$-arbitrarily close to K and $\widetilde{K} \subset \text{Int}(N(K))$. Observe that $N(K)$ is also a closed tubular neighborhood of $\widetilde{K}$. Then, by Theorem 3.3 there exists an isotopic copy $\widetilde{K}$ of the knot K' contained in $\mathcal{S}_k \cap \partial N(K)$ the 1-skeleton of $\partial N(K)$.

In this section, we will prove that there exists a sequence of cubulated moves that carries K into K'.

Lemma 4.4. *There exists a finite sequence of cubulated moves that carries K into K'.*

Proof. Observe that K and K' are isotopic and both are contained into the 1-skeleton of $N(K)$. Next, we will construct an annulus A contained in the 2-skeleton of $\mathcal{N}_k$ such that its boundary will consist of two connected components, namely K and K'.

Let $B = \{Q \subset \mathcal{N}_k \,|\, Q \cap K \neq \emptyset,\ Q \cap K' \neq \emptyset\}$.

Notice that B consists of a finite number of cubes, say m. We will enumerate them in such a way that consecutive numbers belong to neighboring cubes (cubes sharing a common face) and the cube and the m-cube is neighbor of the first one. To construct the annulus in the 2-skeleton of $N(K)$, whose boundary is $K \cup K'$ we will look at all cases of $Q_j \in B$ and find F_j which consists of either one face, two neighboring faces or a face union an edge of Q_j that will form said annulus. The boundary of these F_j's will intersect both K and K', hence faces corresponding to neighboring cubes will share a common edge.

Claim 1: If $Q_j \cap K$, $Q_j \in B$ consists of a vertex and $Q \cap K'$ consists of either a vertex or an edge, then $K \cup K' \subset \cup_{Q_i \in B \setminus \{Q_j\}} Q_i$. In other words, Q_j is superfluos.
Proof of Claim 1. We will look at all cases.

(a) Each intersection ($Q_j \cap K$ and $Q_j \cap K'$) consists of a vertex. This implies that each knot passes through two edges sharing the corresponding vertex and these edges belong to cubes in $B \setminus \{Q_j\}$.

(b) $Q_j \cap K$ consists of a vertex v and $Q_j \cap K'$ consists of an edge e or viceversa (see Figure 1). Suppose that both intersections are contained in the same face $F \subset Q$ (see Figure 1(a)). Notice that K passes through two edges sharing the vertex v. These two edges belong to neighboring cubes of Q_j and K' also intersects them. This implies that one of these two neighboring cubes contains both v and e.
Now, suppose that the vertex v is opposed to e, *i.e.*, there is no edge of Q_j joining v to any of the end-points of e (see Figure 1(b)). As in the previous case, K passes through two edges sharing the vertex v and these two edges belong to neighboring cubes of Q_j. Since K' also intersects them, we have that one of these cubes must contain v and one of the end-points of e. By construction of K', it follows that K' does not intersect any of the other two neighboring cubes of Q_j containing v. Therefore, the claim 1 follows. □



Figure 1: *$Q_j \cap K$ consists of a vertex v and $Q \cap K'$ consists of an edge e.*

For simplicity, we will assume that B does not contained superfluous cubes.

Claim 2: Let $Q_j \in B$. Then $K' \cap Q_j$ can be reduce by face boundary moves, to a path of 3 edges.

Proof of Claim 2. We will consider all possible cases such that $K' \cap Q_j$ consists of at least 4 edges.

Case 1. Suppose that $K \cap Q_j$ consists of a vertex v. Since $K \cap K' = \emptyset$, it follows that $K' \cap Q_j$ can be contained in 1,2 or 3 faces of Q_j and these faces do not contained v. By a combinatorial analysis, we have that $K' \cap Q_j$ is a path of at most 6 edges.

(*a*) $K' \cap Q_j$ consists of six edges. So three edges are contained in a face F_1, two edges in a neighboring face F_2 and one edge in the remaining face F_3 (see Figure 2). In this case, we apply twice the face boundary move to obtain that $K' \cap Q_j$ can be reduce a two edges path.



Figure 2: *$K' \cap Q_j$ consists of a six edges path.*

(*b*) $K' \cap Q$ consists of five edges. In these case, three edges are contained in a face F_1 and two in a neighboring face F_2, then we apply the face boundary move on F_1 to obtain that $K' \cap Q_j$ can be reduce a three edges path (see Figure 3).



Figure 3: *$K' \cap Q_j$ consists of a five edges path.*

(c) $K' \cap Q_j$ consists of four edges. If three edges are contained in a face F_1 and one edge in a neighboring face F_2, then we apply the face boundary move on F_1 to obtain that $K' \cap Q_j$ can be reduce a two edges path (see Figure 4). If two edges are contained in a face F_1 and two edges in a neighboring face F_2, then we apply the face boundary move on F_1 and then apply again the face boundary move on F_2 to obtain that $K' \cap Q_j$ can be reduce a two edges path (see Figure 5).



Figure 4: *$K' \cap Q_j$ consists of a four edges path.*

Case 2. K consists of an edge. Then $K' \cap Q_j$ may be contained in 1 or 2 faces.

(a) Three edges of $K' \cap Q_j$ are contained in a face F_1 and two edges in the neighboring face F_2, then we apply the face boundary move on F_1 to obtain that $K' \cap Q_j$ can be reduce a one edge path (see Figure 6).



Figure 5: $K' \cap Q_j$ *consists of a four edges path.*



Figure 6: $K' \cap Q_j$ *consists of a five edges path.*

(b) Three edges of $K' \cap Q_j$ are contained in a face F_1 and one edge in the neighboring face F_2. Then we apply the face boundary move on F_1 to obtain that $K' \cap Q_j$ can be reduce a two edges path (see Figure 7).



Figure 7: $K' \cap Q_j$ *consists of a four edges path.*

Case 3. K consists of two neighboring edges. In this case, $K' \cap Q_j$ is contained in one face, hence $K' \cap Q_j$ consists of a path of at most three edges and applying the face boundary move, if necessary, we can assume that $K' \cap Q_j$ consists of one or two edges.

This proves claim 2. □

Notice that by claim 2, we will consider that $K' \cap Q_j$ consists of a path of at most three edges. Next, we will construct F_j describing all possible cases of $Q_j \in B$.

Case 1. Suppose that $K \cap Q_j$ consists of a vertex v. By claims 1 and 2, we have that $K' \cap Q_j$ can consist of a path of two or three edges.

(*a*) $K' \cap Q_j$ is a three edges path (see Figure 8). In this case, up to face boundary move, we may assume that v and two neighboring faces (faces sharing a common edge) of $K' \cap Q_j$ are contained in the same face F. Then F_j is the union of F and the remaining edge of $K' \cap Q_j$.



Figure 8: $K \cap Q_j$ *consists of a vertex.*

(*b*) $K' \cap Q_j$ is a two edges path. We have the following arrangements.

1. $K' \cap Q_j$ and v lie on the same face $F \subset Q_j$. Then $F_j = F$. See Figure 9.



Figure 9: $K \cap Q_j$ *and* v *lie on the same face.*

2. $K' \cap Q_j$ and v lie on opposite faces. We have two possibilities that are equivalent via a face boundary move. See



Figure 10: $K \cap Q_j$ *and* $K' \cap Q_j$ *lie on opposite faces.*

Figure 10. By the conditions imposed in B, we have that neighbors cubos of Q_j must intersect K and K', and this can happen only in one case (see Figure 11). Then applying face boundary moves in the neighbors cubes, we see that it is not neccesary to take F_j; *i.e.*, $F_j = \emptyset$.



Figure 11: F_j *is empty.*

3. The remaining configuration appears in Figure 12. By the same argument of claim 1 (b), this configuration is not possible.



Figure 12: *This configuration is not possible.*

Case 2. Suppose that $K \cap Q_j$ consists of an edge. By claim 2, we can assume that $K' \cap Q_j$ consists of one or two edges.

(a) $K' \cap Q_j$ is a three edges path. Since $K \cap K' = \emptyset$, we have that K' must be contained in the two faces of Q_j that do not intersect K. If K' is contained in one face, then applying a face boundary move, we can assume that K' consists of just one edge. Thus, we will consider that K' is contained in two faces. See Figure 13. So F_j is the union of the two faces that intersect both knots.



Figure 13: $K' \cap Q_j$ *consists of a three edges path.*

(b) $K' \cap Q_j$ is a two edges path.

1. Four vertices lie on a face $F \subset Q_j$ and the remaining vertex is joined by an edge e to some vertex of C. Let $F_j = F \cup e$. See Figure 14.



Figure 14: $K \cap Q_j$ *consists of an edge.*

2. The remaining possible configuration appears in Figure 15 and applying a face boundary move is equivalent to a configuration described above (see Figure 14).

(c) $K' \cap Q_j$ is an one edge path.

1. $K' \cap Q_j$ and $K \cap Q_j$ lie on the same face $F \subset Q_j$. Then $F_j = F$ (see Figure 16).



Figure 15: *Equivalent configuration.*



Figure 16: $K' \cap Q_j$ *and* $K \cap Q_j$ *lie on the same face.*

2. $K' \cap Q_j$ and $K \cap Q_j$ lie on opposite edges, *i.e.*, the vertices w_1, $w_2 \in K$ and the vertices v_1, $v_2 \in K'$ satisfy that v_1, w_1 lie on the face $F_1 \subset Q_j$ and v_2, w_2 lie on a opposite face $F_2 \subset Q_j$; in such a way that the edge $\overline{v_1 v_2}$ is parallel to the edge $\overline{w_1 w_2}$. *i.e.* v_1, w_1 are opposite vertices in F_1, and v_2, w_2 are opposite vertices in F_2. See Figure 17.



Figure 17: $K' \cap Q_j$ *and* $K \cap Q_j$ *are opposite edges.*

Observe that there exists a finite sequence of neighbor cubes, $Q_{i+1}, \ldots, Q_j, \ldots, Q_{r-1}$, keeping this arrangement and it changes in the cubes Q_i and Q_r, in such a way that each intersection

$K' \cap Q_i$, $K' \cap Q_r$ consists of two neighboring edges. We can assume, applying induction if necessary, that this happens in the neighbor cubes of Q_j; *i.e.* $i = j-1$ and $r = j+1$. Let $F_1 = Q_{j-1} \cap Q_j$ and $F_2 = Q_{j+1} \cap Q_j$.
Consider the cube Q_{j-1}. Then we have two possible arrangements.

(a) Suppose that its vertices w_1, w_3, v_1, v_3, v satisfy the following: w_1, w_3 belong to K, v_1, v_3, v belong to K', the edge $\overline{v_1 v_3}$ is oppposited to the edge $\overline{w_1 w_3}$ and the vertices v_3, v, w_3 lie on the opposite face of F_1, F_2'. See Figure 18.



Figure 18: *Possible arrangements.*

Thus, the cube Q_{j-1} has the arrangement described in case $2(b)1$. We first apply a face boundary move to Q_{j-1}, so the edges $\overline{vv_3} \cup \overline{v_1 v_3}$ are replaced by the edges $\overline{vz_1} \cup \overline{v_1 z_1}$ obtaining F_{j-1}. Then we have in Q_j the same arrangement in $2(b)1$ so we apply a face boundary move to obtain F_j.

(b) Suppose that $K \cap Q_{j-1} = w_1$. We apply a face boundary move to Q_{j-1} (see case$2(b)1$) in such a way that the edges $\overline{vv_3} \cup \overline{v_1 v_3}$ are replaced by the edges $\overline{vz_1} \cup \overline{v_1 z_1}$. Then there exists another cube $Q \subset A$ such that v, z_1, $w_1 \in Q \cap Q_{j-1}$; hence Q_{j-1} is superfluous. In Q_j we have the case described in $2(b)1$, so we apply a face boundary move to obtain F_j. Notice that $F_{j-1} = \emptyset$.

3. The remaining possible configuration appears in Figure 19 and is very similar to the case $1(b)2$. By the same argument

$F_j = \emptyset$.



Figure 19: $F_j = \emptyset$.

Case 3. Suppose that $K \cap Q_j$ consists of two neighboring edges. Then $K \cap Q_j$ is contained in a face $F \subset A_j$

(a) $K' \cap Q_j$ is a three edges path. Since $K \cap K' = \emptyset$, then K' can be contained in five possible edges, four of them belong to the opposite face of F, say F' and the remaining edge is e. Hence K' consists of e and two edges l_1, $l_2 \subset F'$ (see Figure 20). Applying a face boundary move, if necessary, we can assume that e and l_1 belong to a face F_1 and l_2 and a edge of $K \cap Q_j$ are in a neighboring face F_2. We have that $F_j = F_1 \cup F_2$.



Figure 20: $K \cap Q_j$ *and* $K' \cap Q_j$ *lie on two neighboring faces.*

(b) $K' \cap Q_j$ is a two edges path.

1. $K \cap Q_j$ and $K' \cap Q_j$ lie on two neighboring faces (faces sharing a common edge) of Q_j. Let F_j be the union of these two faces. See Figures 21 and 22.
2. $K' \cap Q_j$ lie on the opposite face F' in such a way that there is only one face F_1 such that both $F_1 \cap K$ and $F_1 \cap K'$ consists on a edge, respectively. Then $F_j = F_1$. See Figure 23



Figure 21: $K \cap Q_j$ *and* $K' \cap Q_j$ *lie on two neighboring faces.*



Figure 22: $K \cap Q_j$ *and* $K' \cap Q_j$ *lie on two neighboring faces.*

Let $\mathcal{A} = \cup F_j$. By the construction, F_j consists of either one face or two neighboring faces, except the case $2(b)1$ where F_j is the union of a face and an edge e . Notice that this edge e is also contained in some other face F_i. In fact, if $e \subset K$ then there exists another cube Q_i containing e, and by hypothesis $Q_i \cap K'$ must contain another edge, hence $e \subset F_i$. If $e \subset K'$ then there exists a neighbor cube Q_r of Q_j such that $Q_r \cap K$ contains an edge, hence $e \subset F_r$. This implies that $\mathcal{A}$ is the union of 2-faces F_j, where F_j is contained into the 2-skeleton of $\mathcal{N}_k$.

Observe also that by construction, the boundary of each F_j is composed by edges belonging to K and K' and two edges e_j and e'_j that do not belong to neither K nor K' and F_j was taken in such a way that the intersection $F_j \cap F_{j+1}$ is one of these two edges e_j, e'_j. Therefore $\mathcal{A}$ is an annulus contained into the 2-skeleton of $\mathcal{N}_k$ whose boundary consists of two connected components, namely K and K'. $\square$

Now, we will carry the knot K' onto the knot K via a finite number of cubulated moves. Notice that $\mathcal{A}$ is the union of m squares which are enumerated; in particular, given the square Q_n, the square Q_{n+1} is a neighboring square of Q_n, *i.e.*, $Q_{n+1} \cap Q_n$ consists of an edge l_n.



Figure 23: *$K \cap Q_j$ and $K' \cap Q_j$ lie on opposite faces.*

We will use induction on m. Consider Q_1. We apply the face boundary move to Q_1 in such a way that the edges belonging to K' replace the others (by construction $Q_1 \cap K = \emptyset$). Next, we consider Q_2. Observe that Q_1 and Q_2 share an edge belonging to K'. Then we apply again a face boundary move, so the edges belonging to K' replace the others. We continue inductively. Notice that if $l \subset K$ then $l \subset \partial \mathcal{A}$, thus l is not a common edge of two squares Q_i and Q_j in $\mathcal{A}$, hence if l is replaced by an edge belonging to K' we have that this replacement will stay in the following steps. Therefore, the result follows. $\square$

4.1.2 Smooth case

Suppose that $\widetilde{K}_1$ and $\widetilde{K}_2$ are isotopic smooth knots in $\mathbb{R}^3$. Let J be the isotopy cylinder of $\widetilde{K}_1$ and $\widetilde{K}_2$. Then J is a smooth submanifold of codimension two in $\mathbb{R}^4$. By Theorem 3.3, there exists an isotopic copy of J, say J', contained in the 2-skeleton of the canonical cubulation $\mathcal{C}$ of $\mathbb{R}^4$. Notice that J' is sliced by connected level sets of p. Moreover there exist integer numbers m_1 and m_2 and cubic knots K'_1 and K'_2 isotopic to $\widetilde{K}_1$ and $\widetilde{K}_2$, respectively; such that $p^{-1}(t) \cap J' = K'_1$ for all $t \leq m_1$ and $p^{-1}(t) \cap J' = K'_2$ for all $t \geq m_2$.

Our goal it to prove that K'_1 is equivalent to K'_2 by cubulated moves,

Let $p : \mathbb{R}^4 \hookrightarrow \mathbb{R}$ be the projection onto the last coordinate. Thus $p^{-1}(t) = \mathbb{R}^3_t$ is an affine hyperplane parallel to the space $\mathbb{R}^3 \times \{0\}$.

Lemma 4.6. *Each hyperplane $\mathbb{R}^3_t$ has a canonical cubulation.*

Proof. Since $\mathbb{R}^3_t$ is an affine hyperplane parallel to the hyperplane $\mathcal{P} = \mathbb{R}^3 \times \{0\} \subset \mathbb{R}^4$ and $\mathcal{P}$ has a canonical cubulation $\mathcal{C}_{\mathcal{P}}$ given by the restriction of the cubulation $\mathcal{C}$ of $\mathbb{R}^4$ to it; *i.e.*, $\mathcal{C}_{\mathcal{P}}$ is the decomposition into cubes which are the images of the unit cube $I^3 = \{(x_1, x_2, x_3, 0) \mid 0 \leq x_i \leq 1\}$ by translations by vectors with integer coefficients. Then $\mathbb{R}^3_t$ has a canonical cubulation coming from the translation of the cubulation $\mathcal{C}_\mathcal{P}$ by the vector $(0, 0, 0, t)$. □

Definition 4.7. A 2-cell (2-face) Q of the canonical cubulation $\mathcal{C}$ of $\mathbb{R}^4$ is called horizontal if $p(Q)$ is a constant in $\mathbb{N}$. A 2-cell is vertical if $p(Q) = [n, n+1]$ for some $n \in \mathbb{N}$.

Definition 4.8. Let Σ be a cubulated surface; *i.e. a surface contained in the 2-skeleton of $\mathcal{C}$*. Let P be a hyperplane in $\mathbb{R}^4$. We say that P and Σ intersect transversally, denoted by $P \pitchfork \Sigma$, if $P \cap \Sigma$ is a polygonal curve.

Lemma 4.9. *Let $\mathbb{R}^3_t$, $t \notin \mathbb{Z}$ be an affine hyperplane. Then $\mathbb{R}^3_t$ intersects J' transversally.*

Proof. Let $\mathbb{P} = \mathbb{R}^3_t$, $t \notin \mathbb{Z}$. By Theorem 3.3, $\mathbb{P} \cap J'$ is connected. Let $x \in \mathbb{P} \cap J'$. Then $x \in Q_i$, where Q_i is a 2-face of the cubulation $\mathcal{C}$. Notice that Q_i is a vertical 2-face, since $t \notin \mathbb{Z}$. So, we have two possibilities: either $x \in \mathrm{Int}(Q_i)$ or x belongs to an edge of Q_i.

1. If $x \in \mathrm{Int}(Q_i)$ then $\mathbb{P} \cap Q_i$ is a linear segment parallel to an edge.
2. If x belongs to an edge of Q_i, then there exists another vertical 2-face Q_j such that $x \in Q_i \cap Q_j$. Thus $\mathbb{P} \cap Q_i$ is a linear segment l_i parallel to an edge, and by the same argument $\mathbb{P} \cap Q_j$ is also a linear segment l_j, and $x \in l_i \cup l_j$.

Therefore, the result follows. □

Corollary 4.10. *For $x \notin \mathbb{Z}$, the set $p^{-1}(x) \cap J'$ is a knot.*

Proof. By the above, $p^{-1}(x) \cap J'$ is a polygonal connected curve. □

Now, for each $n \in \mathbb{N}$ we define

$$K_{-\frac{1}{2}}(n) = p^{-1}(n - \frac{1}{2}) \cap J'$$

and

$$K_{\frac{1}{2}}(n) = p^{-1}(n + \frac{1}{2}) \cap J'.$$

Observe that $K_{-\frac{1}{2}}(n)$ and $K_{\frac{1}{2}}(n)$ are cubic knots.

Let $\mathcal{Q}(\mathcal{C})$ be the set of squares (2-cells) belonging to $\mathcal{C}$. Consider the spaces

$$B_{-\frac{1}{2}}(n) = \cup\{Q \in \mathcal{Q}(\mathcal{C}) \,|\, Q \cap K_{-\frac{1}{2}}(n) \neq \emptyset\},$$

$$B_{\frac{1}{2}}(n) = \cup\{Q \in \mathcal{Q}(\mathcal{C}) \,|\, Q \cap K_{\frac{1}{2}}(n) \neq \emptyset\}$$

and $F_0(n) = p^{-1}(n) \cap J'$. By construction $B_{-\frac{1}{2}}(n) = K_{-\frac{1}{2}}(n) \times [0,1]$ and $B_{\frac{1}{2}}(n) = K_{\frac{1}{2}}(n) \times [0,1]$.

Let $B = B_{-\frac{1}{2}}(n) \cup F_0(n) \cup B_{\frac{1}{2}}(n)$. Notice that $B = \mathrm{Cl}(p^{-1}(n-1, n+1) \cap J')$, where Cl denotes closure.

Lemma 4.11. *The space B is homeomorphic to $\mathbb{S}^1 \times I$.*

Proof. Since B is a compact submanifold of J', then by Lemma 3.3, it is also connected. Now J' is homeomorphic to $\mathbb{S}^1 \times \mathbb{R}$, and $J' - B$ has two connected components, hence the result follows. □

Lemma 4.12. *$F_0(n)$ has the homotopy type of S^1.*

Proof. Consider the set B. By Lemma 4.11, we know that $B \cong \mathbb{S}^1 \times I$, where $K_{-\frac{1}{2}}(n) \times \{0\} \cong \mathbb{S}^1 \times \{0\}$ and $K_{\frac{1}{2}}(n) \times \{0\} \cong \mathbb{S}^1 \times \{1\}$. Hence $\tilde{B} = B/(K_{-\frac{1}{2}}(n) \times \{0\} \cup K_{\frac{1}{2}}(n) \times \{1\})$ is homeomorphic to $\mathbb{S}^2$. By Alexander duality, using reduced homology and cohomology groups, we have that $\tilde{H}_0(\tilde{B} - F_0(n), \mathbb{Z}) = \tilde{H}^1(F_0(n), \mathbb{Z})$. One has that $\tilde{B} - F_0(n)$ has two connected components, so $\tilde{H}_0(\tilde{B} - F_0(n), \mathbb{Z}) \cong \mathbb{Z}$. Since $F_0(n) \subset B \cong \mathbb{S}^1 \times I$, we have that either $\pi_1(F_0(n)) \cong \{0\}$ or $\pi_1(F_0(n)) \cong \mathbb{Z}$, but $H^1(F_0(n), \mathbb{Z}) \cong \mathbb{Z}$, hence $\Pi_1(F_0(n)) \cong \mathbb{Z}$. Therefore, $F_0(n)$ has the homotopy type of S^1. □

Next, we are going to describe the subset $F_0(n)$. Notice that the edges of $F_0(n)$ are of four types, which we will denote by T_1, T_2, T_3 and T_4.

- An edge $l \subset F_0(n)$ belongs to T_1 if $l \subset B_{-\frac{1}{2}}(n)$ but $l \not\subset B_{\frac{1}{2}}(n)$.
- An edge $l \subset F_0(n)$ belongs to T_2 if $l \subset B_{\frac{1}{2}}(n)$ but $l \not\subset B_{-\frac{1}{2}}(n)$.
- An edge $l \subset F_0(n)$ belongs to T_3 if $l \subset B_{-\frac{1}{2}}(n) \cap B_{\frac{1}{2}}(n)$.
- An edge $l \subset F_0(n)$ belongs to T_4 if $l \not\subset B_{\frac{1}{2}}(n)$ and $l \not\subset B_{-\frac{1}{2}}(n)$.

Lemma 4.13. *The space B retracts strongly to $F_0(n)$.*

Proof. Since $B = B_{-\frac{1}{2}}(n) \cup F_0(n) \cup B_{\frac{1}{2}}(n)$ is homeomorphic to $\mathbb{S}^1 \times I$, and $B_{-\frac{1}{2}}(n) = K_{-\frac{1}{2}}(n) \times [0,1]$ and $B_{\frac{1}{2}}(n) = K_{\frac{1}{2}}(n) \times [0,1]$, we have that $B_{-\frac{1}{2}}(n) = K_{-\frac{1}{2}}(n) \times [0,1]$ retracts strongly to $K_{-\frac{1}{2}}(n) \times \{1\}$ and $B_{\frac{1}{2}}(n) = K_{\frac{1}{2}}(n) \times [0,1]$ retracts strongly to $K_{\frac{1}{2}}(n) \times \{0\}$. Now $\partial F_0(n) = K_{-\frac{1}{2}}(n) \times \{1\} \cup K_{\frac{1}{2}}(n) \times \{0\}$. Therefore, the result follows. □

By the above, we have copies of $K_{-\frac{1}{2}}(n)$ and $K_{\frac{1}{2}}(n)$ contained in $\partial F_0(n)$. By abuse of notation we will denote them in the same way. Notice that $K_{-\frac{1}{2}}(n)$ is the union of edges of types T_1 and T_3, and $K_{\frac{1}{2}}(n)$ is the union od edges of types T_2 and T_3.

Lemma 4.14. *There exists a finite sequence of cubulated moves that carries the knot $K_{-\frac{1}{2}}(n)$ into the knot $K_{\frac{1}{2}}(n)$.*

Proof. We will show it by cases.
Case 1. Suppose that $K_{-\frac{1}{2}}(n) = K_{\frac{1}{2}}(n)$. The result is obviously true.

Case 2. Suppose that $K_{-\frac{1}{2}}(n) \cap K_{\frac{1}{2}}(n) = \emptyset$. In other words, $K_{-\frac{1}{2}}(n)$ and $K_{\frac{1}{2}}(n)$ do not have edges of type T_3.
Remember that $F_0(n)$ is a cubulated compact surface whose fundamental group is isomorphic to $\mathbb{Z}$. Thus $\partial F_0(n)$ has two connected component; namely $K_{-\frac{1}{2}}(n)$ and $K_{\frac{1}{2}}(n)$, such that their intersection is empty. Hence $F_0(n)$ is divided by squares (2-faces) belonging to the 2-skeleton of $\mathcal{C}$, whose edges are of any of the types T_1, T_2 and T_4.

Next, we will carry the knot $K_{-\frac{1}{2}}(n)$ onto the knot $K_{\frac{1}{2}}(n)$ via a finite number of cubulated moves; *i.e.* we will carry the edges of type T_1 onto the edges of type T_2. Let Q be a square contained in $F_0(n)$. We can assume, up to $(M1)$-move, that if an edge $l \subset Q$ belongs to T_1, then $Q \cap K_{\frac{1}{2}}(n) = \emptyset$ and $Q \cap K_{-\frac{1}{2}}(n)$ consists of either an edge or two neighboring edges. Analogously, if $l \subset Q$ belongs to T_2, then $Q \cap K_{-\frac{1}{2}}(n) = \emptyset$ and $Q \cap K_{\frac{1}{2}}(n)$ consists of either an edge or two neighboring edges.

Since $F_0(n)$ is compact it is the union of a finite number of squares, say m. We will enumerate them in the following way. The first square Q_1 contains an edge of type T_1, and given the square Q_n, the square Q_{n+1} shares an edge l_n with Q_n and whenever it is possible, we choose Q_{n+1} in such a way that l_n is parallel to l_{n-1}.

We will use induction on m. Consider Q_1. We apply the $(M2)$-move to Q_1 replacing the edges of type T_4 by edges of type T_1. We consider Q_2. Observe that Q_1 and Q_2 share an edge of type T_1. Then we apply again the $(M2)$-move replacing the edges of type T_4 by edges of type T_1. We continue inductively. Notice that if $l \subset F_0(n)$ is an edge of type T_2, then $l \subset \partial F_0(n)$; so l is not a common edge of two squares Q_i and Q_j in $F_0(n)$; hence if l is replaced by an edge of type T_1, then this replacement is not modified on any other step. Therefore, the result follows.

Case 3. Suppose that $K_{-\frac{1}{2}}(n) \cap K_{\frac{1}{2}}(n)$ consists of a finite number of points. The surface $F_0(n)$ consists of connected components C_i, $i = 1, \ldots, r$ such that each C_i is the union of squares $Q_{i_1}, \ldots, Q_{i_{m_i}} \in \mathcal{C}$ and the intersection $C_i \cap C_j$ is either empty or a point belonging to $K_{-\frac{1}{2}}(n) \cap K_{\frac{1}{2}}(n)$. Therefore, we apply the previous argument to each C_i.

Case 4. Suppose that the intersection $K_{-\frac{1}{2}}(n) \cap K_{\frac{1}{2}}(n)$ contains an edge of type T_3.
The surface $F_0(n)$ consists of 2-dimensional connected components C_i, $i = 1, \ldots, r$ and paths γ_{ij}, where each C_i is a union of squares $Q_{i_1}, \ldots, Q_{i_{m_i}} \in \mathcal{C}$, and γ_{ij} is a path (or vertex) joining the component C_i with the component C_j. Observe that if γ_{ij} is a path then it is the union of edges of type T_3, hence $\gamma_{ij} \subset K_{-\frac{1}{2}}(n) \cap K_{\frac{1}{2}}(n)$. Moreover $K_{-\frac{1}{2}}(n) \cap K_{\frac{1}{2}}(n) = \cup \gamma_{ij}$ and $\partial F_0(n) = K_{-\frac{1}{2}}(n) \cup K_{\frac{1}{2}}(n)$.

Since $\Pi_1(F_0(n)) \cong \mathbb{Z}$, then C_i has the has the homotopy type either the circle or the disk. Suppose that C_i has the homotopy type of the circle, then $\partial C_i = \partial F_0(n) = K_{-\frac{1}{2}}(n) \cup K_{\frac{1}{2}}(n)$ contains an edge l of type T_3, but l does not belong to any square Q of $F_0(n)$; so l does not belong to C_i. This is a contradiction, hence C_i has the homotopy type of the disk.

By the above, ∂C_i is homeomorphic to $\mathbb{S}^1$ and consists of edges of type T_1 and T_2. Moreover, ∂C_i consists of two arcs A_1 and A_2, such that A_1 is the union of edges of type T_1 and A_2 is the union of edges of type T_2. Now we apply the argument of the case 2, so A_2 is replaced by A_1. Since we have a finite number of components C_i, the result follows. □

Theorem 4.5. *There exists a finite sequence of cubulated moves that carries* K_1' *into* K_2'. *In other words,* K_1' *is equivalent to* K_2' *by cubulated moves.*

Proof of Theorem 4.5. Recall that there exist integer numbers m_1 and m_2 such that $p^{-1}(t) \cap J' = K_1'$ for all $t \leq m_1$ and $p^{-1}(t) \cap J' = K_2'$ for all $t \geq m_2$. Consider the integer $m_1 + 1$. By l Lemma 4.14 there exists a finite number of cubulated moves that carries the knot K_1' into the knot $K_{\frac{1}{2}}(m_1 + 1)$. We continue inductively, and again by Lemma 4.14 there exists a finite number of cubulated moves that carries the knot $K_{\frac{1}{2}}(m_2 - 1)$ into the knot K_2'. Since, we have a finite number of integers contained in the interval $[m_1, m_2]$, then there exists a finite sequence of cubulated moves that carries K_1' into K_2'. □

5 Discrete knots

In this section we will use the fact that every knot is isotopic to a cubic knot to develop a discrete version of a knot. Since a cubic knot is given by a sequence of edges whose boundaries are in the canonical lattice of points with integer coefficients in $\mathbb{R}^3$, *i*.e., the abelian group $\mathbb{Z}^3$, each knot is determined by a cyclic permutation $(a_1, \ldots, a_n)$ (with some restrictions), $a_i \in \mathbb{Z}^3$. We will describe a discrete version of Seifert surfaces, the fundamental group and the Alexander polynomial of the knots in terms of such cyclic permutations by projecting onto a plane P, such that it is injective when restricted to the $\mathbb{Z}^3$-lattice and the image of the $\mathbb{Z}^3$-lattice, Λ_P, is dense. More precisely, the projection of each knot is determined by a cyclic permutation $(w_1, \ldots, w_n)$ (with some restrictions), $w_i \in \Lambda_P$.

Let $\mathcal{C}$ be the standard cubulation of $\mathbb{R}^3$ and let $\mathcal{S}$ be the corresponding 1-skeleton of $\mathcal{C}$ (scaffolding), which by definition is determined by the vertices in the lattice $\mathbb{Z}^3$ consisting of point with integer coeffcients, and by edges which are contained in straight lines parallel to the coordinate axis and passing through points in the lattice. These straight lines belong to three families of parallel lines: The family $\mathcal{F}_1$ of lines parallel to the x-axis, the family $\mathcal{F}_2$ of lines parallel to the y-axis and the family $\mathcal{F}_3$ of lines parallel to the z-axis.

5.1 Discrete description of cubic knots

Given a cubic knot $K \subset \mathbb{R}^3$, we have that K is a polygonal simple curve whose vertices lie on the lattice $\mathbb{Z}^3$ and whose edges lie on straight lines belonging to the families $\mathcal{F}_i$, $i = 1, 2, 3$ of straight lines parallel to the coordinate axis x_i. This implies that we can write K as a *cyclic permutation* *of points* $(v_1, v_2, \dots, v_n)$ such that $v_i \in \mathbb{Z}^3$, $v_i \neq v_j$ $1 \leq i, j \leq n$ and v_i is joined to v_{i+1} by a unit edge and v_n is likewise joined to v_1 by a unit edge. Observe that the expression $(v_1, v_2, \dots, v_n)$ represents a equivalence class, since $(v_1, v_2, \dots, v_n) = (v_2, v_3, \dots, v_n, v_1) = \dots = (v_n, v_1, \dots, v_{n-1})$. Conversely, a cyclic permutation of points $(v_1, v_2, \dots, v_n)$ satisfying the above conditions determines a unique knot $K \subset \mathcal{S}$. Notice that if K is oriented, we can associate to it a unique cyclic permutation such that the numbering of the $v_i's$ is compatible with this orientation. If K is not oriented there are two cycles that can be associated to K, $(v_1, v_2, \dots, v_n)$ and its inverse $(v_n, v_{n-1}, \dots, v_1)$

Let e_1, e_2, e_3 be the canonical coordinate vectors in $\mathbb{R}^3$. We can also write the cubic knot K as an *"anchored cycle"* permutation of directions $K = \{v_1, (e_{i_1}, \dots, e_{i_n})\}$. In other words, cubic knots can be coded as words over the alphabet $\{e_1^{\pm}, e_2^{\pm}, e_3^{\pm}\}$ plus the initial vertex.

We now can describe the cubulated moves in $\mathbb{R}^3$ using cycle notation.

M1 Subdivision: We subdivide each edge of K by m equal edges. Thus $\{v_1, (e_{i_1}, \dots, e_{i_n})\}$ is equivalent to

$$\{v_1, (\frac{1}{m}e_{i_1}, \dots, \frac{1}{m}e_{i_1}, \dots, \frac{1}{m}e_{i_n}, \dots, \frac{1}{m}e_{i_n})\},$$

where each vector $\frac{1}{m}e_{i_k}$ appears m times (see Figure 24).



Figure 24: *Subdivision move by* $\frac{1}{2}$.

M2 Face boundary move: Let F be a 2-dimensional unit square. Remember that the move M2 consists in replacing the arc $A \subset \partial F$ by the arc $A' = \partial F - A$. If we write K as an anchored cyclic permutation of directions, this move is expressed as follows:

1. If the anchored cyclic permutation of directions is given by

$$\{v_1, (e_{i_1}, \ldots, e_{i_{k-1}}, \mathbf{e_{i_k}}, \mathbf{e_{i_{k+1}}}, -\mathbf{e_{i_k}}, e_{i_{k+3}}, \ldots, , e_{i_n})\}$$

   and $\{v_1, (e_{i_1}, \ldots, e_{i_{k-1}}, \mathbf{e_{i_{k+1}}}, e_{i_{k+3}}, \ldots, e_{i_n})\}$
   are equivalent (see Figure 25).
   In addition, we have:

$$\{\mathbf{v_1}, (\mathbf{e_{i_1}}, e_{i_2} \ldots, e_{i_{n-2}}, -\mathbf{e_{i_1}}, \mathbf{e_{i_n}})\}$$

   and $\{\mathbf{v_1} + \mathbf{e_{i_1}}, (e_2, \ldots, e_{i_{n-2}}, \mathbf{e_{i_n}})\}$ are equivalent, and

$$\{\mathbf{v_1}, (\mathbf{e_{i_1}}, e_{i_2}, e_{i_3}, \ldots, e_{i_{n-1}}, -\mathbf{e_{i_2}})\}$$

   and $\{\mathbf{v_1} + \mathbf{e_{i_2}}, (\mathbf{e_1}, e_{i_3}, \ldots, e_{i_{n-2}}, \mathbf{e_{i_n}})\}$ are equivalent.



Figure 25: *Face boundary move.*

2. If the anchored cyclic permutation of directions is given by

$$\{v_1, (e_{i_1}, \ldots, e_{i_{k-1}}, \mathbf{e_{i_k}}, \mathbf{e_{i_{k+1}}}, e_{i_{k+2}}, \ldots, e_{i_n})\}$$

   and $\{v_1, (e_{i_1}, \ldots, e_{i_{k-1}}, \mathbf{e_{i_{k+1}}}, \mathbf{e_{i_k}}, e_{i_{k+2}}, \ldots, e_{i_n})\}$ are equivalent (see Figure 26).
   In addition:

$$\{\mathbf{v_1}, (\mathbf{e_{i_1}}, e_{i_2} \ldots, e_{i_{n-1}}, \mathbf{e_{i_n}})\}$$

   and $\{\mathbf{v_1} + \mathbf{e_{i_1}} - \mathbf{e_{i_n}}, (e_{i_n}, e_{i_2} \ldots, e_{i_{n-1}}, \mathbf{e_{i_1}})\}$ are equivalent.

Remark 5.1. Note that the face boundary or M2 move in case 2, is only allowed if the vertex that replaces v_{k+1} or v_{i_1}, *i.e.*, $v_{i_{k+1}} + e_{i_{k+1}}$ or $v_1 + e_{i_1} - e_{i_n}$ is not already a vertex of the knot. However, one can always apply the subdivision or M1 move in that case.



Figure 26: *Face boundary move.*

Definition 5.2. Given two anchored cyclic permutation of directions $\{v_1, (e_{i_1}, \dots, e_{i_n})\}$, $\{b_1, (e_{j_1}, \dots, e_{j_r})\}$, we say that they are *equivalent* if there exists a finite sequence of cubulated moves transforming the anchored cyclic permutation of directions $\{h(v_1), (e_{i_1}, \dots, e_{i_n})\}$ into $\{b_1, (e_{j_1}, \dots, e_{j_r})\}$, where h is the translation map given by $h(x) = x + b_1 - v_1$.

By Theorem 1, we know that given two cubic knots K_1 and K_2 in $\mathbb{R}^3$, then they are isotopic if and only if there exists a finite sequence of cubulated moves that carries K_1 into K_2. If we express both knots as anchored cyclic permutations of directions $K_1 = (v_1, e_{i_1}, \dots, e_{i_n})$, $K_2 = (v_2, e_{j_1}, \dots, e_{j_r})$, we have the following

Corollary 5.3. *Two cubic knots K_1 and K_2 in $\mathbb{R}^3$ are isotopic if and only if their anchored cyclic permutations of directions $K_1 = \{v_1, (e_{i_1}, \dots, e_{i_n})\}$ and $K_2 = \{b_1, e_{j_1}, \dots, e_{j_r})\}$ are equivalent.*

5.2 Discrete description of a suitable cubic knot projection

Many invariants of a knot are computed using suitable projections. We will start proving that for any oriented cubic knot $K = (v_1, v_2, \dots, v_n)$, there exists a generic projection p over a suitable fixed plane P. Next, we will represent $\widehat{K} = p(K)$ as a cyclic permutation of points $(w_1, w_2, \dots, w_n)$ and we will describe some invariants of K in terms of the points w_i, $i = 1, \dots, n$.

5.2.1 A canonical generic projection

Lemma 5.4. *There exists a hyperplane P such that the orthogonal projection $p : \mathbb{R}^3 \to P$ is injective in the vertices of $\mathcal{S}$ (the lattice $\mathbb{Z}^3$) and is also injective in each of the families $\mathcal{F}_i$, $i = 1, 2, 3$.*

Proof. Let $N = (1, \pi, \pi^2)$, where π is the well-known transcendental number. Let P be the plane through the origin in $\mathbb{R}^3$ orthogonal to N. Let us show that $p : \mathbb{R}^3 \to P$ is injective when restricted to the lattice $\mathbb{Z}^3$. Let (x_0, y_0, z_0) be a point in P and $L = \{t(1, \pi, \pi^2) + (x_0, y_0, z_0) : t \in \mathbb{R}\}$ the line which is orthogonal to P and passes through (x_0, y_0, z_0). Let us suppose that L contains two points with integer coefficients, $P_1 = (N, M, K) = (x_0 + t_1, y_0 + \pi(t_1), z_0 + \pi^2(t_1))$ and $P_2 = (N', M', K') = (x_0 + t_2, y_0 + \pi(t_2), z_0 + \pi^2(t_2))$. If we consider the point $P_1 - P_2$, which belongs to L, we must have that $P_1 = P_2$ since otherwise π and π^2 would be algebraic numbers which is a contradiction since π is transcendental. Therefore p restricted to the lattice $\mathbb{Z}^3$ is injective. Now we will prove that the projection p restricted to the family $\mathcal{F}_i$, $i = 1, 2, 3$ is injective. First we will consider the case $i = 1$. Let l and l' be two lines belonging to $\mathcal{F}_1$. Suppose that $p(l) = p(l')$. Let Q be the plane in $\mathbb{R}^3$ containing both l and l'. Since p restricted to the coordinate planes is injective, we can assume that Q is not parallel to a coordinate plane. Therefore, a normal vector to Q is of the form $(0, a, b)$ where both a and b are nonzero. Also $p(Q)$ is the line $p(l) = p(l')$. This means that Q contains a translate of the vector $(1, \pi, \pi^2)$ normal to P. That is, $(1, \pi, \pi^2) + x$ belongs to Q, (here x is a point in Q). Hence the interior product of $(1, \pi, \pi^2)$ and $(0, a, b)$ is zero, which is impossible because it would mean that π is a rational multiple of π^2. By a similar argument p restricted to each of the families $\mathcal{F}_2$ and $\mathcal{F}_3$ is also injective. $\square$

Lemma 5.5. *For $p : \mathbb{R}^3 \to P$ defined as above, $p(\mathbb{Z}^3) := \Lambda_P$ is an additive dense subgroup of P. Furthermore, the images $p(\mathcal{F}_i)$ of each of the families $\mathcal{F}_i$, $i = 1, 2, 3$ are dense and invariant under the group of translations of the dense group Λ_P. In addition, for $i = 1, 2, 3$, $p(\mathcal{F}_i)$ is a one dimensional dense subgroup of the additive Lie group $(\mathbb{R}^3, +)$.*

Proof. Let e_1, e_2, e_3 be the canonical coordinate vectors in $\mathbb{R}^3$ and $f_i = p(e_i)$, $i = 1, 2, 3$ be their projections in P. Since p restricted to the coordinate lines is injective, we have that f_1, f_2 and f_3 are pairwise linearly independent. Consider the lattice $\mu = af_1 + bf_2$, $a, b \in \mathbb{Z}$. It is enough to prove that the points kf_3, $k \in \mathbb{Z}$ are dense in the torus P/μ. Since $f_3 = \alpha f_1 + \beta f_2$, α, $\beta \in \mathbb{R}$, we will prove that both α and β are irrational numbers. Suppose that $f_3 = \alpha f_1 + \frac{n}{m} f_2$. Then $p(mf_1 - nf_2 - m\alpha f_3) = 0$, *i.e.*, $(m, -n, -m\alpha) \in \ker(p)$. As $\ker(p)$ is the line orthogonal to P, $L_N = \{t(1, \pi, \pi^2) : t \in \mathbb{R}\}$, it follows that $(m, -n, -m\alpha) = t(1, \pi, \pi^2)$ for some $t \in \mathbb{R}$. This implies that π is rational, which is a contradiction.

Remark 5.6. The set Λ_P is a dense subgroup of rank 3 of P. In fact, if e_1, e_2, e_3 denote the canonical coordinate vectors, then $\mathbb{Z}^3 = \{m_1 e_1 +$

$m_2e_2 + m_3e_3 \,|\, m_1, m_2, m_3 \in \mathbb{Z}\}$ is an additive subgruop of $\mathbb{R}^3$ and since the projection p is a group homomorphism, we have that $\Lambda_P = \{m_1f_1 + m_2f_2 + m_3f_3 \,|\, m_i \in \mathbb{Z};\; f_j = p(e_j)\; j = 1, 2, 3\}$ is a free abelian group with three generators.

Let $K = (v_1, v_2, \ldots, v_n)$ be an oriented cubic knot in $\mathbb{R}^3$. By applying a translation if necessary, we can assume that K is contained in the positive octant of $\mathbb{R}^3$, that is, that all the coordinates of points in K are positive. Let $\widehat{K} = p(K)$ be its projection into the plane P. Thus $\widehat{K}$ is a polygonal curve contained in P with some self-intersections in $\widehat{\mathcal{S}} = p(\mathcal{S})$. Notice that the self-intersections are not contained in Λ_P and by Lemma 5.4, are transverse. This implies that the projection p is regular. The projections of the vertices of K are contained in Λ_P, and are called *vertices* of $\widehat{K}$. The straight line segments joining them are projections of unit edges. The self-intersections of $\widehat{K}$ are called *inessential vertices.*

By the above, we can write $\widehat{K}$ as a *cyclic permutation of points* $(w_1, w_2, \ldots, w_n)$ where $w_i \in \Lambda_P$, $w_i \neq w_j$ $1 \leq i, j < n+1$ and w_i is joined to w_{i+1} by a straight line segment whose preimage is a unit edge, where w_{n+1} is taken to mean w_1 (compare the cubic knot case). In general,

Definition 5.7. An *oriented discrete knot* $\widehat{K}$ is the equivalence class of the n cyclic permutations of n points $(w_1, w_2, \ldots, w_n)$ in $\Lambda_P \subset P$ such that $w_{i+1} - w_i = \pm f_j$, $j \in \{1, 2, 3\}$ and $w_1 - w_n = \pm f_j$.

As in the cubic knot case, we use an alternative description of $\widehat{K}$ as *anchored cyclic permutation* of directions $\{w_1, (f_1, \ldots, f_n)\}$.

Definition 5.8. An inessential vertex (or crossing) c of $\widehat{K}$ is given by $c = \overline{w_kw_{k+1}} \cap \overline{w_jw_{j+1}}$ for $w_k, w_{k+1}, w_j.w_{j+1} \in \widehat{K}$.

Now, we will describe inessential vertices of $\widehat{K}$. For this purpose, let us consider an orthonormal basis of the plane P, $\beta = \{\frac{1}{A}(\pi, -1, 0),\ \frac{1}{B}(\pi^2, \pi^3, -1-\pi^2)\}$ where $A = \sqrt{\pi^2 + 1}$ and $B = \sqrt{2\pi^2 + 2\pi^4 + \pi^6 + 1}$. Now consider four points w_{i_1}, w_{i_2}, w_{i_3} and $w_{i_4} \in \widehat{K}$ whose coordinates in β are $w_{i_j} = (x_j\,,\, y_j)$. We will establish a criteria to know when the line segment $\overline{w_{i_1}w_{i_2}}$ intersects the line segment $\overline{w_{i_3}w_{i_4}}$.

Lemma 5.9. *Let w_{i_1}, w_{i_2}, w_{i_3} and $w_{i_4} \in \widehat{K}$, whose coordinates are $w_{i_j} = (x_j\,,\, y_j)$. Let $u_{r,s} = w_{i_r} - w_{i_s}$ and consider the 2×2 matrices, $A = [u_{2,3}\; u_{4,3}]$, $B = [u_{1,3}\; u_{4,3}]$, $C = [u_{3,1}\; u_{2,1}]$ and $D = [u_{4,1}\; u_{2,1}]$. Then the line segment $\overline{w_{i_1}w_{i_2}}$ intersects the line segment $\overline{w_{i_3}w_{i_4}}$ if and only if* $\det(A)\det(B) < 0$ *and* $\det(C)\det(D) < 0$ *(see Figure 27).*

Proof. See chapter 33 sections 1 and 2 in [4]. □



Figure 27: *Criteria de crossing points.*

Remark 5.10. While applying Lemma 5.9 one needs to consider only the quadruples of points where $i_2 = i_1 + 1$ and $i_4 = i_3 + 1$.

Suppose that the line segment $l_{1,2} = \overline{w_{i_1} w_{i_2}}$ intersects the line segment $l_{3,4} = \overline{w_{i_3} w_{i_4}}$. The next step is to determine which crosses "over" the other. Let w_i, w_{i+1}, w_j and w_{j+1} be vertices such that the line segments $\overline{w_i w_{i+1}}$ and $\overline{w_j w_{j+1}}$ intersect. That is, there is a crossing point between them. The line segments $\overline{w_i w_{i+1}}$ and $\overline{w_j j w_{j+1}}$ are the image under the projection p of two edges of the cubulation which are part of K, and whose end points are v_i, v_{i+1} and v_j, v_{j+1} respectively. We will say $l_{1,2}$ crosses *over* $l_{3,4}$ if the edges $\overline{v_j v_{j+1}}$ is closer to P than $\overline{v_i v_{i+1}}$, and we will say it is an *overcrossing.* In the opposite case we will say it is an *under crossing.*

These edges are parallel to two different canonical coordinate vectors. Let e_s be the third coordinate vector. If we consider the projection onto the s-th coordinate, then $\pi_s(v_i) = \pi_s(v_{i+1}) = M$ and $\pi_s(v_j) = \pi_s(v_{j+1}) = R$. Observe that M and R cannot be equal since this would mean that the edges belong to the same plane, parallel to a coordinate plane, and on these planes the projection is injective.

We claim that the crossing point is an undercrossing if and only if $M < R$. (Recall we are assuming K is in the first quadrant.) Consider the following family of planes $\mathcal{P}$: each plane in $\mathcal{P}$ is parallel to the coordinate plane whose s-th coordinate is equal to zero, and is therefore of the form $P_c = \{(x_1, x_2, x_3) : x_s = c\}$. We consider the planes of this form where

$c \in \mathbb{Z}$. Then the edge $v_i v_{i+1}$ is contained in P_M and the edge $v_j v_{j+1}$ is contained in P_R. Now, any line normal to P intersects P_M before it intersects P_R if and only if $M < R$. $\square$

Suppose that c is a crossing point of the segment $l_j = \overline{w_{j+1} w_j}$ over the segment $l_i = \overline{w_{i+1} w_i}$. Consider the vectors $u_i = w_{i+1} - w_i$, $u_j = w_{j+1} - w_j$ and construct the 2×2-matrix $M = [u_j \; u_i]$, we have two possible configurations. If $\det(M) > 0$, we say that c is a *positive crossing*. If $\det(M) < 0$, c is a *negative crossing* (see Figure 28).



Figure 28: *Positive and negative crossings.*

Examples 5.11. 1. Let K be the trefoil knot. K is represented by a cyclic permutation of points $K = (v_1, v_2, \dots, v_{24})$ (see Figure 29), where

$$\begin{array}{llll}
v_1 = (0,0,0), & v_2 = (0,1,0), & v_3 = (0,1,1), & v_4 = (0,1,2), \\
v_5 = (0,1,3), & v_6 = (0,0,3), & v_7 = (0,-1,3), & v_8 = (1,-1,3), \\
v_9 = (1,-1,2), & v_{10} = (1,-1,1), & v_{11} = (1,0,1), & v_{12} = (1,1,1), \\
v_{13} = (1,2,1), & v_{14} = (1,2,2), & v_{15} = (0,2,2), & v_{16} = (-1,2,2), \\
v_{17} = (-1,1,2), & v_{18} = (-1,0,2), & v_{19} = (0,0,2), & v_{20} = (1,0,2), \\
v_{21} = (2,0,2), & v_{22} = (2,0,1), & v_{23} = (2,0,0), & v_{24} = (1,0,0).
\end{array}$$

Let β be an orthonormal base of the plane P given by two vectors $e_1 = \frac{1}{A}(\pi, -1, 0)$ and $e_2 = \frac{1}{B}(\pi^2, \pi^3, -1-\pi^2)$; where $A = \sqrt{\pi^2+1}$ and $B = \sqrt{2\pi^2 + 2\pi^4 + \pi^6 + 1}$. The projection $\widehat{K}$ of K into P is represented, according to β, by a cyclic permutation of points $\widehat{K} = (w_1, w_2, \dots, w_{24})$, where



Figure 29: *Cubic trefoil knot.*

$$\begin{array}{ll}
w_1 = (0,0), & w_2 = (\frac{-1}{A}, \frac{\pi^3}{B}), \\
w_3 = (\frac{-1}{A}, \frac{\pi^3-\pi^2-1}{B}), & w_4 = (\frac{-1}{A}, \frac{\pi^3-2\pi^2-2}{B}), \\
w_5 = (\frac{-1}{A}, \frac{\pi^3-3\pi^2-3}{B}), & w_6 = (0, \frac{-3\pi^2-3}{B}), \\
w_7 = (\frac{1}{A}, \frac{-\pi^3-3\pi^2-3}{B}), & w_8 = (\frac{\pi+1}{A}, \frac{-\pi^3-2\pi^2-3}{B}), \\
w_9 = (\frac{\pi+1}{A}, \frac{-\pi^3-\pi^2-2}{B}), & w_{10} = (\frac{\pi+1}{A}, \frac{-\pi^3-1}{B}), \\
w_{11} = (\frac{\pi}{A}, \frac{-1}{B}), & w_{12} = (\frac{\pi-1}{A}, \frac{\pi^3-1}{B}),
\end{array}$$

$$\begin{array}{ll}
w_{13} = (\frac{\pi-2}{A}, \frac{2\pi^3-1}{B}), & w_{14} = (\frac{\pi-2}{A}, \frac{2\pi^3-\pi^2-2}{B}), \\
w_{15} = (\frac{-2}{A}, \frac{2\pi^3-2\pi^2-2}{B}), & w_{16} = (\frac{-\pi-2}{A}, \frac{2\pi^3-3\pi^2-2}{B}), \\
w_{17} = (\frac{-\pi-1}{A}, \frac{\pi^3-3\pi^2-2}{B}), & w_{18} = (\frac{-\pi}{A}, \frac{-3\pi^2-2}{B}), \\
w_{19} = (0, \frac{-2\pi^2-2}{B}), & w_{20} = (\frac{\pi}{A}, \frac{-\pi^2-2}{B}), \\
w_{21} = (\frac{2\pi}{A}, \frac{-2}{B}), & w_{22} = (\frac{2\pi}{A}, \frac{\pi^2-1}{B}), \\
w_{23} = (\frac{2\pi}{A}, \frac{2\pi^2}{B}), & w_{24} = (\frac{\pi}{A}, \frac{\pi^2}{B}).
\end{array}$$

In this case, we have three crossings: the segment $\overline{w_6w_5}$ over the segment $\overline{w_{19}w_{18}}$, the segment $\overline{w_{21}w_{20}}$ over the segment $\overline{w_{11}w_{10}}$ and the segment $\overline{w_{12}w_{11}}$ over the segment $\overline{w_1w_{24}}$.

2. Let K be the figure eight knot. So K is represented by a cyclic permutation of points $K = (v_1, v_2, \ldots, v_{24}, v_{40})$ (see Figure 30), where

$$\begin{array}{llll}
v_1 = (1,2,3), & v_2 = (2,2,3), & v_3 = (3,2,3), & v_4 = (4,2,3), \\
v_5 = (4,3,3), & v_6 = (4,4,3), & v_7 = (4,5,3), & v_8 = (4,5,2), \\
v_9 = (4,5,1), & v_{10} = (4,4,1), & v_{11} = (4,3,1), & v_{12} = (4,2,1), \\
v_{13} = (4,1,1), & v_{14} = (3,1,1), & v_{15} = (2,1,1), & v_{16} = (2,1,2), \\
v_{17} = (2,1,3), & v_{18} = (2,1,4), & v_{19} = (2,2,4), & v_{20} = (2,3,4), \\
v_{21} = (2,4,4), & v_{22} = (2,5,4), & v_{23} = (2,6,4), & v_{24} = (2,6,3), \\
v_{25} = (2,6,2), & v_{26} = (3,6,2), & v_{27} = (4,6,2), & v_{28} = (5,6,2), \\
v_{29} = (5,5,2), & v_{30} = (5,4,2), & v_{31} = (4,4,2), & v_{32} = (3,4,2), \\
v_{33} = (3,4,3), & v_{34} = (3,4,4), & v_{35} = (3,4,5), & v_{36} = (2,4,5), \\
v_{37} = (1,4,5), & v_{38} = (1,3,5), & v_{39} = (1,2,5), & v_{40} = (1,2,4),
\end{array}$$



Figure 30: *Cubic eight knot.*

The projection $\widehat{K}$ of K into P is represented, according to β, by a cyclic permutation of points $\widehat{K} = (w_1, w_2, \ldots, w_{40})$, where

$w_1 = (\frac{-2+\pi}{A}, \frac{-3-2\pi^2+2\pi^3}{B}),$ $w_2 = (\frac{-2+2\pi}{A}, \frac{-3-\pi^2+2\pi^3}{B}),$
$w_3 = (\frac{-2+3\pi}{A}, \frac{-3+2\pi^3}{B}),$ $w_4 = (\frac{-2+4\pi}{A}, \frac{-3+\pi^2+2\pi^3}{B}),$
$w_5 = (\frac{-3+4\pi}{A}, \frac{-3+\pi^2+3\pi^3}{B}),$ $w_6 = (\frac{-4+4\pi}{A}, \frac{-3+\pi^2+4\pi^3}{B}),$
$w_7 = (\frac{-5+4\pi}{A}, \frac{-3+\pi^2+5\pi^3}{B}),$ $w_8 = (\frac{-5+4\pi}{A}, \frac{-2+2\pi^2+5\pi^3}{B}),$
$w_9 = (\frac{-5+4\pi}{A}, \frac{-1+3\pi^2+5\pi^3}{B}),$ $w_{10} = (\frac{-4+4\pi}{A}, \frac{-1+3\pi^2+4\pi^3}{B}),$
$w_{11} = (\frac{-3+4\pi}{A}, \frac{-1+3\pi^2+3\pi^3}{B}),$ $w_{12} = (\frac{-2+4\pi}{A}, \frac{-1+3\pi^2+2\pi^3}{B}),$
$w_{13} = (\frac{-1+4\pi}{A}, \frac{-1+3\pi^2+\pi^3}{B}),$ $w_{14} = (\frac{-1+3\pi}{A}, \frac{-1+2\pi^2+\pi^3}{B}),$
$w_{15} = (\frac{-1+2\pi}{A}, \frac{-1+\pi^2+\pi^3}{B}),$ $w_{16} = (\frac{-1+2\pi}{A}, \frac{-2+\pi^3}{B}),$
$w_{17} = (\frac{-1+2\pi}{A}, \frac{-3-\pi^2+\pi^3}{B}),$ $w_{18} = (\frac{-1+2\pi}{A}, \frac{-4-2\pi^2+\pi^3}{B}),$
$w_{19} = (\frac{-2+2\pi}{A}, \frac{-4-2\pi^2+2\pi^3}{B}),$ $w_{20} = (\frac{-3+2\pi}{A}, \frac{-4-2\pi^2+3\pi^3}{B}),$
$w_{21} = (\frac{-4+2\pi}{A}, \frac{-4-2\pi^2+24\pi^3}{B}),$ $w_{22} = (\frac{-5+2\pi}{A}, \frac{-4-2\pi^2+5\pi^3}{B}),$
$w_{23} = (\frac{-6+2\pi}{A}, \frac{-4-2\pi^2+6\pi^3}{B}),$ $w_{24} = (\frac{-6+2\pi}{A}, \frac{-3-\pi^2+6\pi^3}{B}),$
$w_{25} = (\frac{-6+2\pi}{A}, \frac{-2+6\pi^3}{B}),$ $w_{26} = (\frac{-6+3\pi}{A}, \frac{-2+\pi^2+6\pi^3}{B}),$
$w_{27} = (\frac{-6+4\pi}{A}, \frac{-2+42\pi^2+46\pi^3}{B}),$ $w_{28} = (\frac{-6+5\pi}{A}, \frac{-2+3\pi^2+6\pi^3}{B}),$
$w_{29} = (\frac{-5+5\pi}{A}, \frac{-2+3\pi^2+5\pi^3}{B}),$ $w_{30} = (\frac{-4+5\pi}{A}, \frac{-2+3\pi^2+4\pi^3}{B}),$
$w_{31} = (\frac{-4+4\pi}{A}, \frac{-2+2\pi^2+4\pi^3}{B}),$ $w_{32} = (\frac{-4+3\pi}{A}, \frac{-2+\pi^2+4\pi^3}{B}),$
$w_{33} = (\frac{-4+3\pi}{A}, \frac{-3+4\pi^3}{B}),$ $w_{34} = (\frac{-4+3\pi}{A}, \frac{-4-\pi^2+4\pi^3}{B}),$
$w_{35} = (\frac{-4+3\pi}{A}, \frac{-5-2\pi^2+4\pi^3}{B}),$ $w_{36} = (\frac{-4+2\pi}{A}, \frac{-5-3\pi^2+4\pi^3}{B}),$
$w_{37} = (\frac{-4+\pi}{A}, \frac{-5-4\pi^2+4\pi^3}{B}),$ $w_{38} = (\frac{-3+\pi}{A}, \frac{-5-4\pi^2+2\pi^3}{B}),$
$w_{39} = (\frac{-2+\pi}{A}, \frac{-5-4\pi^2+2\pi^3}{B}),$ $w_{40} = (\frac{-2+\pi}{A}, \frac{-4-3\pi^2+2\pi^3}{B}),$

In this case, we have four crossings: the segment $\overline{w_{20}w_{19}}$ over the segment $\overline{w_2w_1}$, the segment $\overline{w_7w_6}$ over the segment $\overline{w_{32}w_{31}}$, the segment $\overline{w_{31}w_{30}}$ over the segment $\overline{w_{11}w_{10}}$, and the segment $\overline{w_{36}w_{35}}$ over the segment $\overline{w_{21}w_{20}}$.

5.2.2 Cubulated moves in P

Let K be a cubic knot and $\widehat{K}$ be its projection. We will describe the "cubulated moves" for $\widehat{K}$. Notice that these moves correspond to the cubulated moves described in Section 5.2.2, hence we use the same term.

Consider $\widehat{K} = (w_1, w_2, \ldots, w_n)$. Notice that $\widehat{K}$ can also be described as an anchored cyclic permutation of directions $\{w_1, (f_1, \ldots, f_n)\}$ where $f_i = w_{i+1} - w_i$, $i = 1, \ldots, n$ and $f_{n+1} = w_1 - w_{n+1}$. We will call w_1 the *anchor* of the knot. Note that applying a cyclic permutation σ to the vertices of $\widehat{K}$ (or the directions of $\widehat{K}$) means that the anchor point changes to $\sigma(w_1)$, where σ is the corresponding permutation on $(w_1, w_2, \ldots, w_n)$.

Now, we will describe the cubulated moves in the plane P using cycle notation.

1. **PM1-move** equivalent to the M1 or subdivision move. The oriented discrete knots $\{w_1, (f_1, \dots, f_n)\}$, and
$$\{w_1, (\frac{1}{m}f_1, \dots, \frac{1}{m}f_1, \dots, \frac{1}{m}f_n, \dots, \frac{1}{m}f_n)\},$$
are equivalent, where each vector $\frac{1}{m}f_i$ appears m-times (see Figure 24).

2. **PM2-move** equivalent to the M2 or face boundary move.

   (a) The oriented discrete knots
   $$\{w_1, (f_1, \dots f_{i-1}, \mathbf{f_i}, \mathbf{f_{i+1}}, \mathbf{f_i^{-1}}, f_{i+3}, \dots, f_n)\}$$
   and $\{w_1, (f_1, \dots f_{i-1}, \mathbf{f_{i+1}}, f_{i+3}, \dots, f_n)\}$ are quivalent (see Figure 25).

   In addition, we have:
   $$\{\mathbf{w_1}, (\mathbf{f_1}, f_2 \dots, f_{n-2}, -\mathbf{f_1}, \mathbf{f_n})\}$$
   and $\{\mathbf{w_1} + \mathbf{f_1}, (f_2, \dots, f_{n-2}, \mathbf{f_n})\}$ are equivalent, and
   $$\{\mathbf{w_1}, (\mathbf{f_1}, f_2, f_3, \dots, f_{n-1}, -\mathbf{f_2})\}$$
   and $\{\mathbf{w_1} + \mathbf{f_2}, (\mathbf{f_1}, f_3, \dots, f_{n-2}, \mathbf{f_n})\}$ are equivalent.

   (b) The oriented discrete knots
   $$\{w_1, (f_1, \dots f_{i-1}, \mathbf{f_i}, \mathbf{f_{i+1}}, f_{i+2}, \dots, f_n)\}$$
   and $\{w_1, (f_1, \dots f_{i-1}, \mathbf{f_{i+1}}, \mathbf{f_i}, f_{i+2}, \dots, f_n)\}$ are equivalent (see Figure 26).

   In addition:
   $$\{\mathbf{w_1}, (\mathbf{f_1}, f_2 \dots, f_{n-1}, \mathbf{f_n})\}$$
   and $\{\mathbf{w_1} + \mathbf{f_1} - \mathbf{f_n}, (f_n, f_2 \dots, f_{n-1}, \mathbf{f_1})\}$ are equivalent.

As in the discrete cubic case, we have that

Definition 5.12. Given two oriented discrete knots $\{w_1, (f_1, \dots, f_n)\}$ and $\{w'_1, (f'_1, \dots, f'_r)\}$, we say that they are equivalent if there exists a finite sequence of cubulated moves which transforms the oriented discrete knot $\{h(w_1), (h(f_1), \dots, h(f_n))\}$ into $\{w'_1, (f'_1, \dots, f'_r)\}$, where h is the translation map given by $h(x) = x + a,\ a \in P$ where $h(w_1) = w'_1$.

As a consequence of Corollary 5.3 and Lemma **??**, it follows that

Corollary 5.13. *Two oriented cubic knots K_1 and K_2 in $\mathbb{R}^3$ are isotopic if and only if their projections, the oriented discrete knots $\widehat{K_1} = \{w_1, (f_1, \dots, f_n)\}$ and $\widehat{K_2} = \{w'_1, (f'_1, \dots, f'_r)\}$ are equivalent.*

5.2.3 Fundamental group

Let $\widehat{K} = (w_1, w_2, \dots, w_n, w_{n+1} = w_1)$ be an oriented discrete knot and $c_1, c_2, \dots, c_r$ be its inessential vertices. We will compute the fundamental group of K, denoted by $\Pi_1(K)$, using the Wirtinger presentation (see [13], [7]). We will start describing the generators of $\Pi_1(K)$.

Suppose that the inessential vertex c_j is the crossing point of the linear segment $\overline{w_{k_j} w_{k_j+1}}$ over the linear segment $\overline{w_{i_j} w_{i_j+1}}$. Now, we are going to rearrange the inessential vertices c_j according to the index i_j in such a way that $i_1 < i_2 < \cdots < i_r$. This allows us to describe $\widehat{K}$ using the cyclic permutation $(c_{i_1}, c_{i_2}, \dots, c_{i_r})$. Let γ_i be the segment of $\widehat{K}$ whose endpoints are c_i and c_{i+1} (where $c_{r+1} = c_1$). Since there is a bijection between the set of segments γ_i, $i = 1, \dots, r$ and the set of generators of $\Pi_1(K)$, we have that the set of generators of $\Pi_1(K)$ is $\{\alpha_1, \dots, \alpha_r\}$.

Next, we will describe the set of relations among the generators. By the Wirtinger presentation, we know that for each inessential vertex c_j corresponds a relation among the three generators α_j, α_{j+1} and α_s, where the index s satisfies that the segment $\overline{w_{k_j} w_{k_j+1}}$ is contained in γ_s. Thus

- If c_j is a positive crossing, then the relation is $\alpha_s \alpha_j = \alpha_{j+1} \alpha_s$.
- If c_j is a negative crossing, then the relation is $\alpha_j \alpha_s = \alpha_s \alpha_{j+1}$.

5.2.4 Seifert surface

We now want to look at the case of Seifert surfaces, developed in [14].

Seifert Theorem: Given an arbitrary oriented knot (or link) K, then there exists in $\mathbb{R}^3$ an orientable, connected surface, F, that has as its boundary K.

To construct F the Seifert algorithm uses an oriented diagram of K. Suppose the diagram has r crossings. Then we will generate (possibly overlapping) disks by *resolving* the crossings, *i.e.*, replacing the crossing by two disjoint arcs that respect the orientation (see Figure 31). We are left with a collection of s simple closed curves called *Seifert curves.* Now we can consider each one of these closed curves to be a boundary of a disk, and the disks can be connected at each crossing by a twisted band, such that the twist in the band corresponds to the crossing (so we need r bands). We thus obtain a *Seifert surface* F for the knot. The genus of F is $\frac{(1-s+r)}{2}$. The *Seifert genus* of a knot is the minimal genus possible for a Seifert surface of that knot. (For more details see [13], [7]).



Figure 31: *Resolving a crossing of a diagram of K.*

We go back to our construction. Let $\widehat{K} = (w_1, w_2, \dots, w_n,)$ be an oriented discrete knot and $c_1, c_2, \dots, c_r$ be its inessential vertices, where $c_i = \overline{w_j w_{j+1}} \cap \overline{w_k w_{k+1}}$. Let $\mathcal{A} = \{(w_j, w_{j+1}),\, (w_k, w_{k+1}) \,|\, i = 1, \dots, r\}$. We define a permutation

$$\sigma : (w_1, w_2, \dots, w_n) \to (w_1, w_2, \dots, w_n)$$

according to the splicing process. In other words, the vertex w_l is sent to w_{l+1} if $(w_l, w_{l+1}) \notin \mathcal{A}$. Otherwise $((w_l, w_{l+1}) \in \mathcal{A})$, there exists an inessential vertex c_i and a pair (w_k, w_{k+1}) such that $c_i = \overline{w_l w_{l+1}} \cap \overline{w_k w_{k+1}}$, then w_l is sent to the vertex w_{k+1}.

This permutation can be expressed as a product of disjoint cycles, where each cycle represents a Seifert curve. Summarizing we have the following:

Theorem 5.14. *Let K be a cubic knot in $\mathbb{R}^3$ and $\widehat{K} = (w_1, w_2, \dots, w_n,)$ be its projection. Let $\sigma : (w_1, w_2, \dots, w_n) \to (w_1, w_2, \dots, w_n)$ be the above permutation. Suppose that σ can be expressed as the product of s disjoint cycles, then the number of Seifert curves of $\widehat{K}$ is s.*

Remark 5.15. By the above, we can compute the genus g of the Seifert surface F, since it can be obtained from the formula $g = \frac{1-s+r}{2}$, discussed above, where s is the number of Seifert curves and r is the number of inessential vertices.

5.2.5 The Alexander polynomial

J.W. Alexander defined a polynomial invariant of an oriented knot (subsequently called the Alexander polynomial) in [1]. To calculate it, one must look at the (oriented) diagram of a knot. Assume the diagram has no trivial loops, and r crossings. In the case where there are no trivial loops, the diagram divides the plane into $r+2$ regions. We create an $r \times (r+2)$ incidence matrix (where the rows represent the r crossings and the columns represents the $r+2$ regions, of course). For entry (c_i, r_j) of the matrix, we will set it equal to 0 if crossing c_i is not adjacent to region r_j. Otherwise, we must set it equal to a value in $\{\pm 1, \pm t\}$, depending on the relationship of the region to the incoming undercrossing. If r_j is on the left and before undercrossing we enter $-t$, if on the left and after it, t, if it is on the right and before undercrossing we enter 1, and in the last case -1. We now make our matrix an $r \times r$ matrix by removing any two columns corresponding to two adjacent regions. We obtain a polynomial by calculating the determinant of this matrix. Of course, we have an ambiguity, since depending on which 2 columns we chose to remove, the polynomial may differ by a factor of $\pm t^n$. So we must then divide by the largest possible power of t. Finally, we multiply by -1 if the constant term is negative. The result is the Alexander polynomial.

Let $\widehat{K} = (w_1, w_2, \dots, w_n,)$ be an oriented discrete knot and $c_1, c_2, \dots, c_r$ be its inessential vertices. We will assume that $\widehat{K}$ does not contain any loops, after applying cubulated moves if necessary. We will describe the regions of the knot diagram in terms of the vertices $w_1, w_2, \dots, w_n$.

Remember that $c_j = \overline{w_{i_j} w_{i_j+1}} \cap \overline{w_{k_j} w_{k_j+1}}$. We associate to each c_j the set S_j in the following way (see Figure 28).

- If c_j is a negative crossing, then $S_j = \{(w_{i_j}, w_{k_j+1}), (w_{k_j+1}, w_{i_j+1}), (w_{i_j+1}, w_{k_j}), (w_{k_j}, w_{i_j})\}$.

- If c_j is a positive crossing, then $S_j = \{(w_{i_j+1}, w_{k_j+1}),\ (w_{k_j+1}, w_{i_j}),\ (w_{i_j}, w_{k_j}),\ (w_{k_j}, w_{i_j+1})\}$.

Let $S = \cup_{j=1}^{r} S_j$. We define the permutation $\tau : S \to S$ as follows: $\tau(w_x, w_y) = (w_{\phi(y)}, w_z)$ where $\phi(y)$ is the smallest integer greater or equal to y such that $(w_{\phi(y)}, w_z) \in S \setminus S_j$.

This permutation can be expressed as a product of disjoint cycles. Observe that each cycle determines a circle given by the concatenation of the corresponding vertices where each vertex is taken once. Then there exists a natural bijection between these circles and the regions of the knot diagram. Hence we can compute the Alexander polynomial applying the above procedure.

References

[1] Alexander, J. W. *Topological invariants of knots and links.* Trans. Amer. Math. Soc. (1928) 30 (2): 275306.

[2] E. Artin. *Zur Isotopie zweidimensionalen Flächen im R_4.* Abh. Math. Sem. Univ. Hamburg (1926), 174–177.

[3] Boege, M., Hinojosa, G. and Verjovsky, A. *Any smooth knot $\mathbb{S}^n \hookrightarrow \mathbb{R}^{n+2}$ is isotopic to a cubic knot contained in the canonical scaffolding of* $\mathbb{R}^{n+2}$. Rev. Mat Complutense (2011) 24: 1–13. DOI 10.1007/s13163-010-0037-4.

[4] T. H. Cormen, C. E. Leiserson, R. L. Rivest, C. Stein. *Introduction to Algorithms.* The MIT Press. Third Edition.

[5] N. P. Dolbilin, M. A. Shtan ko, M. I. Shtogrin. *Cubic manifolds in lattices.* Izv. Ross. Akad. Nauk Ser. Mat. 58 (1994), no. 2, 93-107; translation in Russian Acad. Sci. Izv. Math. 44 (1995), no. 2, 301-313.

[6] A. Douady. *Arrondissement des arêtes.* (French) 1961/1962 Séminaire Henri Cartan, 1961/62, Exp. 3 25 pp. Secrétariat mathématique, Paris.

[7] R. H. Fox. *A Quick Trip Through Knot Theory.* Topology of 3-Manifolds and Related Topics. Prentice-Hall, Inc., 1962.

[8] L. Funar. *Cubulations, immersions, mappability and a problem of Habegger.* Ann. scient. Éc. Norm. Sup., 4e série, t. 32, 1999, pp. 681–700.

[9] B. Mazur. *The definition of equivalence of combinatorial imbeddings.* Publications mathématiques de l'I.H.É.S., tome 3 (1959) p.5–17.

[10] S. Matveev, M. Polyak. *Finite-Type Invariants of Cubic Complexes.* Acta Applicandae Mathematicae 75, pp. 125–132, 2003.

[11] J. Milnor. *Morse Theory.* Princeton University Press, 1969.

[12] R. Palais. *Local triviality of the restriction map for embeddings.* Comment. Math. Helv. 34, 1960, 305–312.

[13] D. Rolfsen. *Knots and Links.* Publish or Perish, Inc. 1976.

[14] Seifert, H. *Über das Geschlecht von Knoten.* Math. Annalen (1934) 110 (1): 571–592.

[15] Živaljević, Rade T. *Combinatorial groupoids, cubical complexes, and the Lovász conjecture.* Discrete Comput. Geom. 41 (2009), no. 1, 135–161.

G. Hinojosa. `Facultad de Ciencias`. Universidad Autónoma del Estado de Morelos. Av. Universidad 1001, Col. Chamilpa. Cuernavaca, Morelos, México, 62209.
E-mail address: gabriela@uaem.mx

A. Verjovsky. `Instituto de Matemáticas, Unidad Cuernavaca`. Universidad Nacional Autónoma de México. Av. Universidad s/n, Col. Lomas de Chamilpa. Cuernavaca, Morelos, México, 62209.
E-mail address: alberto@matcuer.unam.mx

C. Verjovsky Marcotte. `Mathematics Department`. St. Edwards University. 3001 South Congress Avenue, Austin, TX 78704, USA.
E-mail address: cynthiavm@gmail.com